

**Spray Recommendations for Downy Mildew**  
 Mary K. Hausbeck, Professor and Extension Specialist  
 Department of Plant Pathology, Michigan State University

**CUCUMBER**

APPLIED BEFORE DISEASE (7-day intervals)	APPLIED AFTER DISEASE (5-day intervals)
<ul style="list-style-type: none"> <li>• Gavel 75WG (5 day PHI)</li> <li>• Presidio 4FL (2 day PHI)</li> <li>• Previcur Flex 6SC (2 day PHI)</li> <li>• Ranman 3.6SC (0 day PHI)</li> <li>• Tanos 50WG (3 day PHI)</li> </ul>	<ul style="list-style-type: none"> <li>• Previcur Flex 6SC (2 day PHI)</li> <li>• Presidio 4FL (2 day PHI)</li> <li>• Ranman 3.6SC (0 day PHI)</li> <li>• Tanos 50WG (3 day PHI)</li> </ul>
Alternate products and mix each with either: <ul style="list-style-type: none"> <li>• Dithane (mancozeb) 3 lb or</li> <li>• Bravo (chlorothalonil) 1.5 pt</li> </ul>	Alternate products and mix each with either: <ul style="list-style-type: none"> <li>• Dithane (mancozeb) 3 lb or</li> <li>• Bravo (chlorothalonil) 2 pt</li> </ul>

**VINE CROPS, such as zucchini, squash, pumpkin, melon and gourds**

APPLIED BEFORE DISEASE (7- to 10-day intervals)	APPLIED AFTER DISEASE (7-day intervals)
<ul style="list-style-type: none"> <li>• Gavel 75WG (5 day PHI)</li> <li>• Presidio 4FL (2 day PHI)</li> <li>• Previcur Flex 6SC (2 day PHI)</li> <li>• Ranman 3.6SC (0 day PHI)</li> <li>• Tanos 50WG (3 day PHI)</li> </ul>	<ul style="list-style-type: none"> <li>• Previcur Flex 6SC (2 day PHI)</li> <li>• Presidio 4FL (2 day PHI)</li> <li>• Ranman 3.6SC (0 day PHI)</li> <li>• Tanos 50WG (3 day PHI)</li> </ul>
Alternate products and mix each with either: <ul style="list-style-type: none"> <li>• Dithane (mancozeb) 3 lb or</li> <li>• Bravo (chlorothalonil) 1.5 pt</li> </ul>	Alternate products and mix each with either: <ul style="list-style-type: none"> <li>• Dithane (mancozeb) 3 lb or</li> <li>• Bravo (chlorothalonil) 2 pt</li> </ul>

[Please note: Gavel 75WG and Dithane are not registered on some vine crops.]